

A PROPERTY OF ALTERNATING GROUPS

HENRY CEJTIN AND IGOR RIVIN

ABSTRACT. We describe an efficient algorithm to write any element of the alternating group A_n as a product of two n -cycles (in particular, we give a simple proof that any element of A_n can be so written). An easy corollary is that every element of A_n is a commutator in S_n .

Consider the alternating group A_n . We will show the following

Theorem 1. *There is an $O(n)$ algorithm to write any element $\sigma \in A_n$ as a product of two n -cycles.*

That any element of A_n can be so written is a result of E. A. Bertram [Bertram72], but his argument is less direct and hence does not seem to lead to an optimal algorithm. On the other hand, Bertram's result is more general.

A corollary (Corr. 7), it is seen that every element of A_n is a commutator in S_n – the well-known fact that A_n is the commutator subgroup of S_n gives a much weaker statement. As an immediate corollary of the (one line) proof of Corr. 7 we obtain an $O(n)$ algorithm to represent any element of A_n as a commutator.

In the sequel we use exclusively the cycle notation for permutations. Both the existence and the algorithmic aspects of the cycle decomposition are discussed in [Knuth68, Section 1.3.3].

In order to prove Theorem 1 we will need a few lemmas:

Lemma 2. *Let $\sigma_1 = (a_1, \dots, a_s)$, and $\sigma_2 = (b_1, \dots, b_t)$. Furthermore suppose that $\{a_1, \dots, a_s\} \cap \{b_1, \dots, b_t\} = \emptyset$. Let $\rho = (a_s, b_t)$. Then $\tau_1 = \sigma_1 \sigma_2 \rho$ is an $s + t$ -cycle, as is $\tau_2 = \rho \sigma_1 \sigma_2$.*

Proof. Since $\tau_2 = \rho \tau_1 \rho^{-1}$, it is enough to show that τ_1 is an $s + t$ -cycle. Since the cycle decomposition is conjugacy invariant, we can assume without loss of generality that

$$\begin{aligned}\sigma_1 &= (1, 2, \dots, s), \\ \sigma_2 &= (s + 1, \dots, s + t), \\ \rho &= (s, s + t).\end{aligned}$$

Date: today.

Then

$$\tau_1(x) = \begin{cases} x+1, & x < s-1, \\ s+t, & x = s-1, \\ s+1, & x = s+t, \\ x+1, & s < x < s+t-1, \\ s, & x = s+t-1, \\ x, & x > s+t \end{cases}$$

The assertion of the lemma follows by inspection. $\square$

Lemma 3. *Suppose $\sigma \in S_n$, where $n = t + s$, can be written as $\sigma_1\sigma_2$, where σ_1 only acts non-trivially on $\{1, \dots, t\}$ and σ_2 only acts non-trivially on $\{t+1, \dots, t+s\}$. Suppose further that σ_1 is a product of two t -cycles and σ_2 is a product of two s -cycles. Then σ is a product of two n -cycles.*

Proof. Let $\tau = (t, t+s)$. Let $\sigma_1 = \rho_{11}\rho_{12}$, while $\sigma_2 = \rho_{21}\rho_{22}$, where the ρ_{ij} are the t - and s -cycles as per hypothesis of the Lemma. Then:

$$\sigma_1\sigma_2 = \rho_{11}\rho_{12}\rho_{21}\rho_{22} = \rho_{11}\rho_{21}\rho_{12}\rho_{22} = (\rho_{11}\rho_{21}\tau)(\tau\rho_{12}\rho_{22}).$$

By Lemma 2 both the parenthesized terms are n -cycles, and the result follows. $\square$

Lemma 4. *Let n be odd. Then any n -cycle ρ is a product of two n -cycles.*

Proof. The cyclic subgroup C_ρ generated by ρ is of order n , $m = \frac{n+1}{2}$ is an integer, and clearly $(\rho^m)^2 = \rho$. Since $n+1$ is relatively prime to n , ρ^m is a generator of C_ρ , hence an n -cycle. $\square$

Lemma 5. *Let $n = 4m$, and let $\sigma \in S_n$ be the product of ρ_1 and ρ_2 , where ρ_1 and ρ_2 are $2m$ -cycles, and ρ_1 and ρ_2 are disjoint. Then σ is a product of two n -cycles.*

Proof. Without loss of generality, we can assume that

$$\begin{aligned} \rho_1 &= (1, 3, \dots, 4m-1), \\ \rho_2 &= (2, 4, \dots, 4m). \end{aligned}$$

It is easy to see that $\sigma = \rho^2$, where

$$\rho = (1, 2, \dots, 4m)$$

$\square$

Lemma 6. *Let $n = 2s + 2t$, where $s < t$. Let $\sigma \in S_n$ be the product of ρ_1 and ρ_2 , where ρ_1 is a $2s$ -cycle and ρ_2 is a $2t$ -cycle, and ρ_1 is disjoint from ρ_2 . Then σ is a product of two n -cycles.*

Proof. First, without loss of generality, assume that

$$\begin{aligned}\rho_1 &= (1, 3, \dots, 4s - 1), \\ \rho_2 &= (2, 4, 6, \dots, 2s + 2t, 4s + 1, 4s + 3, \dots, 2s + 2t - 1).\end{aligned}$$

Now, define $\lambda_1, \lambda_2 \in S_n$ as follows:

$$\lambda_1(x) = \begin{cases} 4s + 1 & x = 1, \\ 1 & x = 4s, \\ 2 & x = 2s + 2t, \\ x + 1 & \text{otherwise.} \end{cases}$$

In other words, λ_1 is the n -cycle

$$(1, 4s + 1, 4s + 2, \dots, 2s + 2t, 2, 3, \dots, 4s).$$

$$\lambda_2(x) = \begin{cases} 1 & x = n, \\ x + 1 & \text{otherwise.} \end{cases}$$

In other words, λ_2 is the n -cycle

$$(1, 2, 3, \dots, n).$$

Now, let us compute $\lambda_1\lambda_2$:

$$\lambda_1\lambda_2(n) = \lambda_1(1) = 4s + 1.$$

If $x \neq n$,

$$\lambda_1\lambda_2(x) = \lambda_1(x + 1) = \begin{cases} 1 & x = 4s - 1, \\ 2 & = n - 1, \\ x + 2 & \text{otherwise.} \end{cases}$$

It follows immediately that

$$\lambda_1\lambda_2 = \rho_1\rho_2.$$

□

Proof of Theorem 1. The proof will proceed by induction on n . If $n = 1$, the result is obvious. Now, consider $\sigma \in A_n$. If $\sigma = \sigma_1\sigma_2$, where σ_1 and σ_2 are even and disjoint, the assertion of Theorem 1 will follow from the induction hypothesis and Lemma 3.

Consider now the cycle decomposition of σ – when we count the cycles below, we *do* count trivial (one-element) cycles.

- **The cycle decomposition of σ has exactly one cycle.** In this case, the statement of the Theorem follows from Lemma 4.
- **The cycle decomposition of σ has exactly two cycles.** The statement of the Theorem follows from Lemmas 5 and 6.

- **The cycle decomposition of σ has more than two cycles.** If one of the cycles is of odd length, then it is an even permutation, and the result follows from the discussion at the beginning of the proof. If no cycle is of odd length, then, since σ is an even permutation, there is an even number of cycles in the cycle decomposition of σ , so

$$\sigma = \rho_1 \rho_2 \dots \rho_{2k} = (\rho_1 \rho_2)(\rho_3 \dots \rho_{2k}).$$

Each of the parenthesized terms is an even permutation, and so σ is a product of two n -cycles by the discussion at the beginning of this proof. □

Corollary 7. *Every element of A_n is a commutator of a pair of elements in S_n .*

Proof. All n -cycles are conjugate, and if ρ is an n -cycle, so is ρ^{-1} . Now let $\sigma \in A_n$. By Theorem 1, $\sigma = \rho_1 \rho_2$, where ρ_1, ρ_2 are n -cycles. Since $\rho_2 = \tau \rho_2^{-1} \tau^{-1}$, we see that $\sigma = [\rho_1, \tau]$. □

REFERENCES

- [Bertram72] E. A. Bertram. *Even permutations as a product of two conjugate cycles*, J. Combinatorial Theory Ser. A **12**(1972), pp. 368-380.
- [Knuth68] D. E. Knuth. *The Art of Computer Programming*, vol. 1 1968, Addison-Wesley Publishing.

SOURCELIGHT TECHNOLOGIES, EVANSTON, IL
E-mail address: `henry@sourcelight.com`

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, PHILADELPHIA
Current address: Mathematics Department, Princeton University
E-mail address: `rivin@math.temple.edu`